



Published in final edited form as:

Pediatrics. 2010 July ; 126(1): 115–128. doi:10.1542/peds.2009-1381.

Respiratory Morbidity and Lung Function in Preterm Infants of 32 to 36 Weeks' Gestational Age

Andrew A. Colin, MD^a, Cynthia McEvoy, MD, MCR^b, and Robert G. Castile, MD^c

^aDivision of Pediatric Pulmonology, Miller School of Medicine, University of Miami, Miami, Florida

^bDivision of Neonatology, Oregon Health & Science University, Portland, Oregon

^cCenter for Perinatal Research, Research Institute at Nationwide Children's Hospital, Ohio State University School of Medicine and Public Health, Columbus, Ohio

Abstract

Normal lung development follows a series of orchestrated events. Premature birth interrupts normal in utero lung development, which results in significant alterations in lung function and physiology. Increasingly, there are reports documenting the broad range of complications experienced by infants aged 34 to 36 weeks' gestational age (GA). Our objective was to summarize the evidence demonstrating respiratory system vulnerability in infants aged 34 to 36 weeks' GA and to review the developmental and physiologic principles that underlie this vulnerability. A comprehensive search for studies that reported epidemiologic data and respiratory morbidity was conducted on the PubMed, Medline, Ovid Biosis, and Embase databases from 2000 to 2009 by using medical subject headings “morbidity in late preterm infants,” “preterm infants and lung development,” “prematurity and morbidity,” and “prematurity and lung development.” Because the number of studies exclusive to infants aged 34 to 36 weeks' GA was limited, selected studies also included infants aged 32 to 36 weeks' GA. Of the 24 studies identified, 16 were retrospective population-based cohort studies; 8 studies were observational. These studies consistently revealed that infants born at 32 to 36 weeks' GA, including infants of 34 to 36 weeks' GA, experience substantial respiratory morbidity compared with term infants. Levels of morbidity were, at times, comparable to those observed in very preterm infants. The developmental and physiologic mechanisms that underlie the increased morbidity rate and alterations in respiratory function are discussed. We also present evidence to demonstrate that the immaturity of the respiratory system of infants 34 to 36 weeks' GA at birth results in increased morbidity in infancy and leads to deficits in lung function that may persist into adulthood.

Keywords

airway conductance; airway patency; bronchopulmonary dysplasia; functional residual capacity; late preterm; respiratory morbidity; tethering

The frequency of late-preterm (34^w–36^w weeks' gestational age [GA]) births has generally increased in the United States over the past 2 decades, with a slight decline in 2007 (10.61% in 1990, 12.80% in 2006, and 12.66% in 2007).¹ Infants of 34 to 36 weeks' GA comprise ~71% of all preterm births (9.03% overall in 2007 [~390 000 live births]).¹ Late-preterm

infants are often considered to be as physiologically and metabolically mature as term infants and, therefore, at low risk for morbidity and mortality.^{2,3}

A modest but growing body of literature has documented the broad range of complications that late-preterm infants may experience. Compared with term infants, late-preterm infants are at higher risk of developing medical complications and have higher rates of mortality during infancy.^{4–8} Despite significant declines in the infant mortality rate over the last decade, the infant mortality rate for late-preterm infants has remained three- to fivefold higher than that for term infants.^{4–7,9} Each weekly reduction in estimated GA at birth was found to be associated with an increased risk of death.¹⁰ Late-preterm infants also face increased morbidity before hospital discharge^{4,11,12} and higher rates of hospital readmission in the first months of life.^{5,11–13} They are 4 times more likely than term infants to have ≥ 1 medical condition diagnosed and 3.5 times more likely to have ≥ 2 conditions diagnosed.^{13,14} Late-preterm infants are often viewed as normal and are discharged early (<2-night hospital stay), which may be a factor in the increased overall risk of rehospitalization.^{12,15}

In addition to numerous neonatal complications, reviewed by Engle et al¹⁴ and Makari and Groothuis,¹⁶ a high incidence of respiratory distress has been reported as one of the most common adverse outcomes among late-preterm infants.^{7,17–21} There has been little information on the long-term consequences of preterm birth and essentially none for late-preterm birth beyond the first year of life. Results of a recent review on the respiratory consequences of preterm birth suggested that many adverse respiratory consequences of preterm birth are likely the result of persistent pulmonary structural abnormalities.²²

Differences in respiratory development between term and late-preterm infants may be illustrated by the susceptibility of preterm infants (33–36 weeks' GA) to infection by such pathogens as respiratory syncytial virus (RSV), for which infection rates among preterm infants (33–36 weeks' GA) more resembled those of younger preterm infants (<33 weeks' GA) than those of term infants.²³

The original purpose of this review was to summarize the recent literature regarding the increased respiratory morbidity in preterm infants aged 34 to 36 weeks' GA, a population that largely presents with a low incidence of surfactant-deficiency–related neonatal respiratory problems. We then provide a physiologic explanation as to why these apparently healthy infants have a high incidence of respiratory problems compared with term infants. The physiologic and developmental characteristics underpinning the immediate vulnerability of this population are described along with emerging evidence of long-term sequelae associated with birth during the saccular stage of lung development. Additional background is presented to provide context for the data summary and possible explanation of the results. This review is intended to inform providers who make decisions regarding delivery of late-preterm infants and who are involved in their care in early life of the higher risks faced by these children.

The American Academy of Pediatrics and the American College of Obstetricians and Gynecologists have defined late-preterm infants as those born between $34\frac{0}{7}$ and $36\frac{6}{7}$ weeks' GA.²⁴ As stated above, our original intent was to focus on this narrow subpopulation; however, for the purposes of this review, we have included studies for which outcomes in infants born between 32 and 36 weeks' GA were reported, because many of the publications of interest often included patient populations a few weeks below the 34- to 36-weeks' GA window.

Online databases (PubMed, Medline, Ovid Biosis, Embase) from 2000 to 2009 were searched by using medical subject headings “morbidity in late preterm infants” and “preterm

infants and lung development” to identify studies that included infants of 34 to 36 weeks' GA and reported ≥ 1 objective quantified outcome (including respiratory morbidity) without regard for sample size. Additional Embase searches included Emtree Thesaurus and free terms “prematurity and morbidity” and “prematurity and lung development.” The articles selected were limited to those published during or after the year 2000 to capture studies that examined lung morbidity such as bronchopulmonary dysplasia (BPD) using the more recent definition from a workshop sponsored by the Eunice Kennedy Shriver National Institute of Child Health and Human Development, the National Heart, Lung, and Blood Institute, and the Office of Rare Diseases.²⁵ The primary objectives, study design and methods, comparison populations, pertinent results (including morbidity and mortality rates of appropriate studies), and study limitations were summarized. Twenty-four studies were identified for inclusion in this review. A summary of the salient features of these studies is provided in Table 1. To describe the underlying respiratory physiology, additional publications are cited that were not limited to the years 2000–2009.

GENERAL MORBIDITY AND MORTALITY

A recent study²⁶ that involved 377 638 term and 26 170 late-preterm (34- to 36-week-GA) infants revealed that 22.2% of the late-preterm infants versus 3.0% of the term infants had newborn morbidity during their birth hospitalization. Preterm infants born at 35 and 36 weeks' GA had 10-fold (relative risk [RR]: 10.2 [95% confidence interval [CI]: 9.7–10.8) and fivefold (RR: 4.8 [95% CI: 4.6–5.1]) higher risk of morbidity, respectively, compared with infants born at 40 weeks' GA. Infants born at 34 weeks' GA had a 20-fold (RR: 20.6 [95% CI: 19.7–21.6]) higher risk of morbidity than infants born at 40 weeks' GA. Relative morbidity increased approximately twofold with every week of decreasing GA earlier than 38 weeks' GA (Fig 1).²⁶ In a retrospective study, McIntire and Leveno⁷ showed neonatal mortality rates of 1.1, 1.5, and 0.5 per 1000 live births at 34, 35, and 36 weeks' GA, respectively, versus 0.2 per 1000 live births at 39 weeks' GA ($P < .001$). Pulver et al⁸ and Santos et al⁵ showed that 34- to 36-week-GA infants were ~7 and 3 times more likely to die in their first month compared with 39- to 42-week-GA and 37- to 41-week-GA infants, respectively. Two retrospective analyses of records of infants born to US residents revealed higher infant mortality rates in 34- to 36-week-GA infants compared with term infants (37–41 and 40 weeks' GA, respectively).^{6,10}

GENERAL RESPIRATORY MORBIDITY

All forms of respiratory morbidity, including transient tachypnea of the newborn, respiratory distress syndrome, pneumonia, and pulmonary hypertension, affect late-preterm infants at a higher rate than infants of more advanced GA.^{7,13,17,19,20} A study by De Luca et al²⁷ revealed a strong age-related trend independent of delivery mode in respiratory morbidity and a >10-fold increase in respiratory morbidity in infants of 34 weeks' GA compared with term infants. A retrospective study in Canada by Kitsomart et al²⁸ revealed significantly worse respiratory outcomes (including prevalence of pneumothorax, rates of positive pressure therapy, and mechanical ventilation assistance; all $P < .001$) in 1481 infants of 34 to 36 weeks' GA compared with 9332 infants of ≥ 37 weeks' GA.

INFECTIOUS RESPIRATORY MORBIDITY

Very preterm infants (<32 weeks' GA), infants with BPD or congenital heart disease, and preterm infants born at 32 to 36 weeks' GA are particularly vulnerable to developing severe lower respiratory tract infections that require frequent hospitalizations.^{13,19,29–32} A study by Boyce et al²³ revealed that infants born at 33 to 36 weeks' GA had similar rates of admission to the hospital for RSV infection as infants born at <33 weeks' GA, and there were substantially higher rates of admission in these 2 populations compared with those in term

infants. Infants born at 33 to 35 weeks' GA had hospital outcomes that were similar to or worse than those of infants born at ≤ 32 weeks' GA regardless of whether RSV infection was confirmed³³ or the infants were hospitalized for nonspecific bronchiolitis.³⁴ After hospitalization for RSV, infants born at 32 to 36 weeks' GA experienced rehospitalization rates twice as high, hospital stays 3 times as long, and outpatient visits twice as frequent as infants of similar GA who were not hospitalized for RSV.³⁵ Prematurity was also associated with an increased risk of bronchiolitis-associated death. The odds ratio for death related to RSV in 32- to 35-week-GA infants compared with term infants was ~ 5 .³⁶

BASIC MECHANISMS OF LUNG DEVELOPMENT

To understand the mechanisms that possibly explain the morbidity in infants born at 34 to 36 weeks' GA, it is necessary to understand lung physiology at this stage of their development.

Stages of Lung Development

Normal lung development is characterized by a carefully choreographed series of prenatal and postnatal events that can be compromised by a variety of factors.³⁷ Early growth and development of the human lung is a continuous process that is highly variable between individuals but, nonetheless, has traditionally been divided into 5 stages. The first is the embryonic phase (26 days to 6 weeks' GA), followed by the pseudoglandular (6–16 weeks' GA), canalicular (16–28 weeks' GA), sacular (28–36 weeks' GA), and alveolar (36 weeks' GA to term) phases. This final phase continues into childhood.^{37,38} Interference with this stepwise process of lung development during any of these phases may render the lung less effective as a gas exchanger and more susceptible to disease.³⁹

The formation of conducting airways and terminal bronchioles occurs during the canalicular period and establishes the platform for gas exchange.³⁷ Bronchial branching to roughly the 16th generation of the bronchial tree is complete by 16 weeks' GA.^{40,41} The 28- to 36-week-GA period, termed the sacular period, is a transitional phase before full maturation of alveoli occurs characterized by an increased number of saccules, primitive alveoli that become gradually more effective as gas exchangers and may be sufficient in number and quality to sustain life in the preterm infant. The alveolar walls in the sacular stage are more compact and thick than the final thin walls of alveoli. They also include a double capillary structure that is reduced to a single one in the mature alveolus. However, the blood vessels are well oriented to the epithelium and protrude into air spaces, forming many thin air-blood interfaces that are capable of carrying out the function of gas exchange that fully matures in the alveolar phase.

Although alveoli may form during the sacular phase, mature alveoli are not uniformly present until 36 weeks' GA.³⁸ During the alveolar phase, the epithelium and interstitium decrease in thickness, air-space walls proliferate, and the capillary network matures to its final single-capillary network.^{38,42} Blood-vessel development, which begins at the earliest stages, continues throughout lung development.⁴³ It is important to note that these structural changes not only affect gas exchange but have profound effects on the mechanical properties of the lung and, hence, the respiratory system as a whole. As noted, premature birth during this critical period may result in significant alteration in lung function and physiology.³⁸

Stable Functional Residual Capacity and Effectiveness of Gas Exchange

Maintenance of a stable and adequate functional residual capacity (FRC) is important for securing stable gas exchange. FRC is determined by the balance between the opposing forces of the chest wall and lung and, thus, is a direct function of their respective mechanical properties. In early life, a compliant chest wall offers little outward recoil to the respiratory system; thus, the elastic characteristics of the respiratory system approximate those of the

lung. The lung is also more compliant (ie, has less elastance) in premature and newborn infants. The lung becomes less compliant (ie, increases in elastance) as it undergoes alveolarization, and the interstitium becomes more intricately woven.⁴⁴ Compliance of the chest wall declines over the first 2 years of life.^{45–48} In early life, the lung–chest-wall equilibrium results in a mechanically determined FRC that is low relative to older children and adults and is an important determinant of age-related vulnerability to hypoxia. To circumvent this limitation, infants actively elevate their FRC by using laryngeal braking during tidal expiration and by initiating inspiration at an end-expiratory volume above that determined by the mechanical properties of the chest wall and lung.^{49,50} An additional mechanism is persistence of inspiratory muscle activity into the expiratory phase, thereby modulating the expiratory flow.⁵¹

The transition from an actively maintained FRC to one that, as in adults and older children, is mechanically determined occurs in term infants late in the first year and into the second year of life.⁵² The timing of this transition coincides with the declining compliance (stiffening) of the chest wall and its gradually increasing contribution to the overall stabilization of the respiratory system (Fig 2).^{46,47} Chest-wall compliance is particularly elevated in premature infants, with the slope of change toward reduced compliance being the steepest between 28 and 40 weeks' GA compared with that at any stage in postnatal life (Fig 2A).⁴⁶ Thus, it is reasonable to assume that breathing with an overly compliant chest wall, high lung compliance, and reduced number of air-containing units is a challenge for infants delivered before term. The challenge of maintaining an FRC that permits stable gas exchange is likely compounded in the premature infant by apneic events, which have been shown to drive the system to critically low lung volumes and result in rapid desaturation.⁵⁰

Airway Tethering

An additional crucial mechanism that secures airway patency and, thus, adequate maintenance of FRC is airway-tethering.^{53–56} Tethering is mediated through the elastic components in alveolar walls that surround bronchi. These elastic fibers are anchored to each other, creating an extended mesh that exerts a circumferential pull on the intraparenchymal airways (Fig 3). This complex elastic network transmits tension from the pleural surface to the individual bronchi. The tension on the system increases during inspiration, resulting in increased airway caliber. The cross-sectional area of the airway decreases with decline in lung volume. Thus, tethering couples lung-volume changes to airway caliber. Tethering of airways is less effective in infants born prematurely, because alveolarization and the associated development of the parenchymal elastic network are still in the sacular stage of development at 32 to 36 weeks' GA. The effect of reduced tethering is decreased airway stability, increased tendency to closure, increased airway resistance, and, ultimately, a tendency to collapse alveolar units in the lung periphery.^{53–56}

Changes in Lung Volume During the Last Trimester of Gestation

Total lung volume undergoes rapid changes during the last trimester of gestation. Calculations by Langston et al³⁸ revealed that at 30 weeks' GA, the lung volume is only 34% of the ultimate lung volume at mature birth, and at 34 weeks only reaches 47% of the final volume at maturity. In contrast, the air-space walls decrease in thickness such that at 30 and 34 weeks, they are 164% (28 μm) and 135% (23 μm), respectively, relative to the ultimate wall thickness at mature birth (17 μm). In parallel, dramatic increases in air-space surface area occur. Surface area increases from 1.0 to 2.0 m^2 at 30 to 32 weeks' GA and to 3.0 to 4.0 m^2 at term. These volume changes likely have direct mechanical implications in reducing the vulnerability caused by a low and unstable FRC. Maturation of the alveolar network improves parenchymal elastance and, therefore, airway-tethering.

ASSOCIATION OF PRETERM (30- TO 36-WEEK-GA) BIRTH WITHOUT CLINICAL LUNG DISEASE WITH ALTERED LUNG DEVELOPMENT AND FUNCTION

Lung development occurs mostly in utero. At term, the lung is in its final stage of development: the alveolar stage.⁵⁷ Normal in utero lung development occurs according to a highly programmed sequence in a stable milieu, one that is significantly more hypoxic relative to the atmosphere (at a fraction of inspired oxygen of 2%–3% at 10–12 weeks' GA, which rises to 8%–10% thereafter). This hypoxic environment represents the norm for lung organogenesis, including vascular development.^{58–60} Early events of trophoblast differentiation are oxygen regulated.⁶¹ It is now recognized that postnatal hyperoxia plays a key role in the development of BPD.⁶² Premature birth interrupts normal in utero lung development and results in an early transition from the hypoxic intrauterine environment to a comparatively hyperoxic atmospheric environment. An inhaled oxygen concentration of 21% represents significant hyperoxia for the preterm infant. Although this relative hyperoxia has not been directly demonstrated to be an independent cause of altered lung maturation, it may play a role in the subsequent development of chronic lung dysfunction.

There is increasing evidence to support the hypothesis that preterm delivery, even in the absence of any neonatal respiratory disease, may have adverse effects on subsequent lung growth and development and that these alterations may persist and worsen during the first 5 years of life.^{63–68} Results of a limited number of studies have shown that premature infants of varying GAs, but born without clinical lung disease, have altered pulmonary function. Reduced airway function in the absence of neonatal respiratory disease was demonstrated at 1 year of age in a population of healthy infants born at 29 to 36 weeks' GA.⁶⁶ Mansell et al reported lower airway conductances and maximum expiratory flows in a group of 5- to 7-year-old children who had been born prematurely but without respiratory problems, which suggests that airway dysfunction persists into childhood.⁶⁹ These effects may not be dissimilar to those experienced by very preterm infants (<32 weeks' GA), who have generally been the focus of attention because of the severity of their respiratory problems immediately after birth and frequent development of BPD.^{70,71} A direct association between premature birth and reduced expiratory flows was demonstrated recently by using the raised-volume rapid-thoracic-compression technique.⁷² In these studies, healthy preterm infants (30–34 weeks' GA [mean: 33.4 weeks' GA]) studied at a mean corrected age of 8 weeks had reduced airway flows in the presence of normal forced vital capacity (FVC) compared with term infants. In a follow-up analysis, the reduced flows did not normalize in these children by 16 months of age, thus demonstrating a lack of “catch-up” growth in airway function by early in the second year of life (Fig 4).⁷³ This result led the authors to conclude that preterm birth was associated with altered lung development. These longitudinal studies that used the raised-volume method confirmed the results of previous studies that suggested a similar conclusion.²²

The long-term significance of reduced airway function early in life was recently emphasized in a longitudinal study that involved a large group of nonselected (enrolled at birth without any specific criteria) infants who had participated in the Tucson Children's Respiratory Study.⁷⁴ The study's results showed that infants whose pulmonary function was in the lowest quartile also had pulmonary function in the lowest quartile through the years of follow-up until early adulthood.⁷⁴ These findings in a normal unselected population suggest that level of pulmonary function in early life tracks and changes little with growth. Data from Weiss and Ware⁷⁵ have suggested that deficits in lung function during early life, especially if associated with lower respiratory illnesses, increase the risk of chronic obstructive

pulmonary disease later in adult life. Of particular importance in this context is the role played by RSV, which is known to affect most children during their first year of life.³²

DISCUSSION

Late-preterm infants (34⁰/₇–36⁶/₇ weeks' GA) account for nearly three-quarters of all preterm births in the United States. However, until recently, the emphasis of most research has been on the very preterm infant (<32 weeks' GA) or the very low birth weight infant (≤1500 g). In this review we focused on respiratory disorders, the predominant consequence, but not the only system affected by, preterm birth (34–36 weeks' GA).^{7,18}

A search of the literature from 2000 to 2009 identified 24 studies that reported respiratory morbidity and mortality in populations that included late-preterm infants, 9 of which specifically addressed the impact of RSV bronchiolitis in this population. The evidence presented in the reviewed articles demonstrates that the respiratory vulnerability of preterm infants born at 32 to 36 weeks' GA, usually considered to be at low risk for subsequent respiratory problems, is more similar to that of very preterm infants (<32 weeks' GA) than of term infants. Consequently, these infants are at high risk of developing medical complications that result in higher rates of respiratory morbidity and mortality compared with term infants, as demonstrated by their higher rates of hospital readmission shortly after discharge and during the following 6 months, longer hospital stays, and increased health care use compared with term infants.^{35,36}

Taken together, the results of the published studies suggest that birth at 32 to 36 weeks' GA is a very important risk factor for respiratory morbidity during infancy and early childhood. In this review, we made an attempt to describe physiologic and developmental phenomena that underlie the vulnerability of infants to respiratory diseases. These mechanisms play a role in all infants but are likely to be of particular importance in the preterm population. The conclusions that we draw regarding late-preterm infant lung function are largely extrapolated from physiologic principles and a developmental continuum between very preterm and term infants. Infants born at 30 to 34 weeks' GA without clinical lung disease have altered lung function that persists throughout infancy.^{72,73} The physiologic deficiencies that result from incomplete lung development described above are likely to account for the early morbidity and vulnerability to infection. With growth in the first to second years of life, the physiologic instability of the chest wall and the inadequate maintenance of lung volume likely correct themselves and should not account for the long-term deficiencies. Therefore, it is apparent that an actual impairment of the potential for full growth is related to early birth. The observation of a direct association between premature birth and reduced forced expiratory flows⁷² suggests that interruption of lung development is an important determinant of the subsequent increase in respiratory morbidity. Deficits in lung function during early life seem to persist until early adulthood⁷⁴ and, if associated with lower respiratory illnesses, may increase the risk of chronic obstructive pulmonary disease later in life.⁷⁵ These findings may have major implications for obstetric and neonatal care.

One of the consequences of altered lung development in preterm infants is their increased vulnerability to RSV infection. Published data have suggested that preterm birth at 33 to 35 weeks' GA significantly increases the risk for severe outcomes among infants who are hospitalized for RSV.³³ RSV hospitalization in healthy premature (32- to 35-week-GA) infants is associated with a significant increase in subsequent health care resource use.³⁵ Welliver et al⁷⁶ proposed that the pathogenesis of viral lower respiratory tract infection is primarily related to the failure to develop an adaptive cytotoxic T-lymphocyte response and that clearance of the virus depends on inefficient innate immune responses mediated by macrophages and neutrophils. The increased vulnerability to RSV in infants born at 32 to 36

weeks' GA suggests that these maturation-dependent deficiencies may be more pronounced in this age group.

The main limitation of the published retrospective analyses is the fact that information on morbidity and mortality was obtained, in most cases, by review of hospital discharge records linked to birth and death certificates, which may have introduced errors in the estimation of GAs and in the accuracy of reporting. Also, the quality of care provided to preterm infants born at 32 to 36 weeks' GA could not be retrospectively evaluated, which made it difficult to compare data between the different studies. Because only published studies were examined, the presence of publication bias cannot be excluded. Despite these limitations, perinatal consequences of late-preterm birth in relation to morbidity and hospitalization ultimately translated into a several-fold greater risk of neonatal mortality among late-preterm infants born at 34 to 36 weeks' GA (1.5–7.9 deaths per 1000 live births) compared with term infants (0.2–2.4 deaths per 1000 live births).^{4–7,9}

Those involved in the care of late-preterm infants should recognize that these infants have, on average, lower airway function than term infants and require diligent evaluation and monitoring, during the neonatal period and throughout early childhood.^{77,78}

The various physiologic studies cited in this review included infants with a spectrum of GAs that spanned prematurity from as early as 30 to 32 weeks' GA, with 34 to 36 weeks' GA representing the upper range of study subjects. Although the demarcation that separated the latter age group is blurred, we think that the conclusions from these studies apply to the older GAs within the study population. Statements regarding long-term and short-term lung function have to be viewed with these limitations in mind.

Substantial research efforts are needed to move toward a more scientific approach in this population. Research attention should be directed to the newborn group without BPD, and within this group, particular emphasis should be placed on late-preterm infants (born at 34–36 weeks' GA). Data from Friedrich et al^{72,73} reveal lung-function deficits in prematurely born infants without BPD and require confirmation in larger populations. Although it seems likely, it has not yet been demonstrated that the subset of those infants with the lowest lung function is responsible for the majority of the observed later increase in morbidity in this population. The lung function and clinical characteristics of the subset of infants at greatest risk for subsequent morbidity and rehospitalization in the first years of life need to be prospectively characterized. Identifying the characteristics of the group of infants at highest risk for later morbidity will enable the development of trials of therapeutic interventions intended to prevent late respiratory morbidity in these infants. Monoclonal antiviral medications have been shown to be effective in reducing late respiratory morbidity in infants born at ≤ 32 weeks' GA and infants with BPD. Definition of a target group of 32- to 36-week-GA infants with high rates of rehospitalization resulting from respiratory illnesses would make therapeutic trials of these agents feasible. It will also be important to better characterize the nature and distribution of the lung abnormalities in these at-risk infants by using additional functional measures (plethysmographic fractional lung volumes, lung-clearance index, diffusion capacity) and lung-imaging techniques. The reviewed studies cited herein focused on morbidities and mortality that occur shortly after birth or during the ensuing 6 months. They did not address whether the clinical outcomes they evaluated had long-term implications. Physiologic studies of the at-risk cohort should be extended to childhood and beyond to investigate the long-term physiologic consequences of late-preterm birth.

Acknowledgments

We acknowledge Physicians World, a division of KnowledgePoint360 Group, LLC, for providing writing and editing support. Additional editorial support was provided by John E. Fincke, PhD, and Gerard P. Johnson, PhD, of Complete Healthcare Communications, Inc. Both were funded by MedImmune.

FINANCIAL DISCLOSURE: Dr Colin has served on advisory boards conducted by MedImmune and serves on Merck's speaker's bureau; Dr McEvoy has served on advisory boards for MedImmune and owns stock in pharmaceutical companies, and MedImmune has provided grant support for her investigator-initiated research; and Dr Castile and Nationwide Children's Hospital receive royalties from nSpire Health, Inc in relation to sales of the Infant Pulmonary Laboratory, and Dr Castile has also served on an advisory board conducted by MedImmune.

ABBREVIATIONS

GA	gestational age
RSV	respiratory syncytial virus
BPD	bronchopulmonary dysplasia
RR	relative risk
CI	confidence interval
FRC	functional residual capacity
FVC	forced vital capacity

REFERENCES

1. Hamilton BE, Martin JA, Ventura SJ. Births: preliminary data for 2007. *Natl Vital Stat Rep* 2009;57(12):1–23. [PubMed: 19754006]
2. American College of Obstetricians and Gynecologists. ACOG educational bulletin. Assessment of fetal lung maturity. Number 230, November 1996. Committee on Educational Bulletins of the American College of Obstetricians and Gynecologists. *Int J Gynaecol Obstet* 1997;56(2):191–198. [PubMed: 9061400]
3. American College of Obstetricians and Gynecologists. ACOG committee opinion: antenatal corticosteroid therapy for fetal maturation. *Obstet Gynecol* 2002;99(5 pt 1):871–873. [PubMed: 11978304]
4. Kramer MS, Demissie K, Yang H, Platt RW, Sauvé R, Liston R. The contribution of mild and moderate preterm birth to infant mortality. Fetal and Infant Health Study Group of the Canadian Perinatal Surveillance System. *JAMA* 2000;284(7):843–849. [PubMed: 10938173]
5. Santos IS, Matijasevich A, Silveira MF, et al. Associated factors and consequences of late preterm births: results from the 2004 Pelotas birth cohort. *Paediatr Perinat Epidemiol* 2008;22(4):350–359. [PubMed: 18578748]
6. Tomashek KM, Shapiro-Mendoza CK, Davidoff MJ, Petrini JR. Differences in mortality between late-preterm and term singleton infants in the United States, 1995–2002. *J Pediatr* 2007;151(5):450–456. e456.e1. [PubMed: 17961684]
7. McIntire DD, Leveno KJ. Neonatal mortality and morbidity rates in late preterm births compared with births at term. *Obstet Gynecol* 2008;111(1):35–41. [PubMed: 18165390]
8. Pulver LS, Guest-Warnick G, Stoddard GJ, Byington CL, Young PC. Weight for gestational age affects the mortality of late preterm infants. *Pediatrics* 2009;123(6) Available at: www.pediatrics.org/cgi/content/full/123/6/e1072.
9. Escobar GJ, Clark RH, Greene JD. Short-term outcomes of infants born at 35 and 36 weeks gestation: we need to ask more questions. *Semin Perinatol* 2006;30(1):28–33. [PubMed: 16549211]
10. Young PC, Glasgow TS, Li X, Guest-Warnick G, Stoddard G. Mortality of late-preterm (near-term) newborns in Utah. *Pediatrics* 2007;119(3) Available at: www.pediatrics.org/cgi/content/full/119/3/e659.

11. Shapiro-Mendoza CK, Tomashek KM, Kotelchuck M, Barfield W, Weiss J, Evans S. Risk factors for neonatal morbidity and mortality among “healthy,” late preterm newborns. *Semin Perinatol* 2006;30(2):54–60. [PubMed: 16731277]
12. Tomashek KM, Shapiro-Mendoza CK, Weiss J, et al. Early discharge among late preterm and term newborns and risk of neonatal morbidity. *Semin Perinatol* 2006;30(2):61–68. [PubMed: 16731278]
13. Wang ML, Dorer DJ, Fleming MP, Catlin EA. Clinical outcomes of near-term infants. *Pediatrics* 2004;114(2):372–376. [PubMed: 15286219]
14. Engle WA, Tomashek KM, Wallman C. “Late-preterm” infants: a population at risk [published correction appears in *Pediatrics*. 2008;121(2):451]. *Pediatrics* 2007;120(6):1390–1401. [PubMed: 18055691]
15. Escobar GJ, Gonzales VM, Armstrong MA, Folck BF, Xiong B, Newman TB. Rehospitalization for neonatal dehydration: a nested case-control study. *Arch Pediatr Adolesc Med* 2002;156(2):155–161. [PubMed: 11814377]
16. Makari D, Groothuis J. Health risks of the late-preterm infant. *Neonatal Today* 2009;4(1):1–8.
17. Clark RH. The epidemiology of respiratory failure in neonates born at an estimated gestational age of 34 weeks or more. *J Perinatol* 2005;25(4):251–257. [PubMed: 15605071]
18. Escobar GJ, Greene JD, Hulac P, et al. Rehospitalisation after birth hospitalisation: patterns among infants of all gestations. *Arch Dis Child* 2005;90(2):125–131. [PubMed: 15665162]
19. Khashu M, Narayanan M, Bhargava S, Osioviich H. Perinatal outcomes associated with preterm birth at 33 to 36 weeks' gestation: a population-based cohort study. *Pediatrics* 2009;123(1):109–113. [PubMed: 19117868]
20. Dani C, Corsini I, Piergentili L, Bertini G, Pratesi S, Rubaltelli FF. Neonatal morbidity in late preterm and term infants in the nursery of a tertiary hospital. *Acta Paediatr* 2009;98(11):1841–1843. [PubMed: 19604170]
21. Melamed N, Klinger G, Tenenbaum-Gavish K, et al. Short-term neonatal outcome in low-risk, spontaneous, singleton, late preterm deliveries. *Obstet Gynecol* 2009;114(2 pt 1):253–260. [PubMed: 19622985]
22. Moss TJ. Respiratory consequences of preterm birth. *Clin Exp Pharmacol Physiol* 2006;33(3):280–284. [PubMed: 16487275]
23. Boyce TG, Mellen BG, Mitchel EF Jr, Wright PF, Griffin MR. Rates of hospitalization for respiratory syncytial virus infection among children in Medicaid. *J Pediatr* 2000;137(6):865–870. [PubMed: 11113845]
24. American Academy of Pediatrics. American College of Obstetricians and Gynecologists. Guidelines for Perinatal Care. 5th ed.. American Academy of Pediatrics; Elk Grove Village, IL: 2005.
25. Jobe AH, Bancalari E. Bronchopulmonary dysplasia. *Am J Respir Crit Care Med* 2001;163(7):1723–1729. [PubMed: 11401896]
26. Shapiro-Mendoza CK, Tomashek KM, Kotelchuck M, et al. Effect of late-preterm birth and maternal medical conditions on newborn morbidity risk. *Pediatrics* 2008;121(2) Available at: www.pediatrics.org/cgi/content/full/121/2/e223.
27. De Luca R, Boulvain M, Irion O, Berner M, Pfister RE. Incidence of early neonatal mortality and morbidity after late-preterm and term cesarean delivery. *Pediatrics* 2009;123(6) Available at: www.pediatrics.org/cgi/content/full/123/6/e1064.
28. Kitsommart R, Janes M, Mahajan V, et al. Outcomes of late-preterm infants: a retrospective, single-center, Canadian study. *Clin Pediatr (Phila)* 2009;48(8):844–850. [PubMed: 19596865]
29. Carbonell-Estrany X, Quero J, Bustos G, et al. Rehospitalization because of respiratory syncytial virus infection in premature infants younger than 33 weeks of gestation: a prospective study. *IRIS Study Group. Pediatr Infect Dis J* 2000;19(7):592–597. [PubMed: 10917214]
30. Law BJ, Langley JM, Allen U, et al. The Pediatric Investigators Collaborative Network on Infections in Canada study of predictors of hospitalization for respiratory syncytial virus infection for infants born at 33 through 35 completed weeks of gestation. *Pediatr Infect Dis J* 2004;23(9):806–814. [PubMed: 15361717]

31. Figueras-Aloy J, Carbonell-Estrany X, Quero J. Case-control study of the risk factors linked to respiratory syncytial virus infection requiring hospitalization in premature infants born at a gestational age of 33–35 weeks in Spain. *Pediatr Infect Dis J* 2004;23(9):815–820. [PubMed: 15361718]
32. Leader S, Kohlhase K. Recent trends in severe respiratory syncytial virus (RSV) among US infants, 1997 to 2000. *J Pediatr* 2003;143(5 suppl):S127–S132. [PubMed: 14615711]
33. Horn SD, Smout RJ. Effect of prematurity on respiratory syncytial virus hospital resource use and outcomes. *J Pediatr* 2003;143(5 suppl):S133–S141. [PubMed: 14615712]
34. Willson DF, Landrigan CP, Horn SD, Smout RJ. Complications in infants hospitalized for bronchiolitis or respiratory syncytial virus pneumonia. *J Pediatr* 2003;143(5 suppl):S142–S149. [PubMed: 14615713]
35. Sampalis JS. Morbidity and mortality after RSV-associated hospitalizations among premature Canadian infants. *J Pediatr* 2003;143(5 suppl):S150–S156. [PubMed: 14615714]
36. Holman RC, Shay DK, Curns AT, Lingappa JR, Anderson LJ. Risk factors for bronchiolitis-associated deaths among infants in the United States. *Pediatr Infect Dis J* 2003;22(6):483–490. [PubMed: 12799502]
37. Joshi S, Kotecha S. Lung growth and development. *Early Hum Dev* 2007;83(12):789–794. [PubMed: 17905543]
38. Langston C, Kida K, Reed M, Thurlbeck WM. Human lung growth in late gestation and in the neonate. *Am Rev Respir Dis* 1984;129(4):607–613. [PubMed: 6538770]
39. Maritz GS, Morley CJ, Harding R. Early developmental origins of impaired lung structure and function. *Early Hum Dev* 2005;81(9):763–771. [PubMed: 16081227]
40. Bucher U, Reid L. Development of the intra-segmental bronchial tree: the pattern of branching and development of cartilage at various stages of intra-uterine life. *Thorax* 1961;16:207–218. [PubMed: 13874265]
41. Kotecha S. Lung growth for beginners. *Paediatr Respir Rev* 2000;1(4):308–313. [PubMed: 16263457]
42. Copland I, Post M. Lung development and fetal lung growth. *Paediatr Respir Rev* 2004;5(suppl A):S259–S264. [PubMed: 14980282]
43. Stenmark KR, Gebb SA. Lung vascular development: breathing new life into an old problem. *Am J Respir Cell Mol Biol* 2003;28(2):133–137. [PubMed: 12540479]
44. Bryan, CA.; Wohl, MEB. Respiratory mechanics in children. In: Macklem, PT.; Mead, J., editors. *Handbook of Physiology: The Respiratory System*. Vol. Vol 3. American Physiological Society; Bethesda, MD: 1986. p. 183-191. Pt 2
45. Davis GM, Coates AL, Papageorgiou A, Bureau MA. Direct measurement of static chest wall compliance in animal and human neonates. *J Appl Physiol* 1988;65(3):1093–1098. [PubMed: 3182479]
46. Gerhardt T, Bancalari E. Chestwall compliance in full-term and premature infants. *Acta Paediatr Scand* 1980;69(3):359–364. [PubMed: 7376862]
47. Papastamelos C, Panitch HB, England SE, Allen JL. Developmental changes in chest wall compliance in infancy and early childhood. *J Appl Physiol* 1995;78(1):179–184. [PubMed: 7713809]
48. Richards CC, Bachman L. Lung and chest wall compliance of apneic paralyzed infants. *J Clin Invest* 1961;40:273–278. [PubMed: 13741257]
49. Kosch PC, Hutchinson AA, Wozniak JA, Carlo WA, Stark AR. Posterior cricoarytenoid and diaphragm activities during tidal breathing in neonates. *J Appl Physiol* 1988;64(5):1968–1978. [PubMed: 3391897]
50. Kosch PC, Stark AR. Dynamic maintenance of end-expiratory lung volume in full-term infants. *J Appl Physiol* 1984;57(4):1126–1133. [PubMed: 6501029]
51. Mortola JP, Milic-Emili J, Noworaj A, Smith B, Fox G, Weeks S. Muscle pressure and flow during expiration in infants. *Am Rev Respir Dis* 1984;129(1):49–53. [PubMed: 6703485]
52. Colin AA, Wohl ME, Mead J, Ratjen FA, Glass G, Stark AR. Transition from dynamically maintained to relaxed end-expiratory volume in human infants. *J Appl Physiol* 1989;67(5):2107–2111. [PubMed: 2600036]

53. Gomes RF, Shardonofsky F, Eidelman DH, Bates JH. Respiratory mechanics and lung development in the rat from early age to adulthood. *J Appl Physiol* 2001;90(5):1631–1638. [PubMed: 11299249]
54. Henschen M, Stocks J, Brookes I, Frey U. New aspects of airway mechanics in preterm infants. *Eur Respir J* 2006;27(5):913–920. [PubMed: 16455837]
55. Moreno RH, Hogg JC, Pare PD. Mechanics of airway narrowing. *Am Rev Respir Dis* 1986;133(6):1171–1180. [PubMed: 3717766]
56. Plopper CG, Nishio SJ, Schelegle ES. Tethering tracheobronchial airways within the lungs. *Am J Respir Crit Care Med* 2003;167(1):2–3. [PubMed: 12502468]
57. Burri, PH. Development and growth of the human lung. In: Macklem, PT.; Mead, J., editors. *Handbook of Physiology: The Respiratory System*. Vol. Vol 3. American Physiological Society; Bethesda, MD: 1997. p. 1-46.Pt 1
58. Burton GJ, Jauniaux E, Watson AL. Maternal arterial connections to the placental intervillous space during the first trimester of human pregnancy: the Boyd collection revisited. *Am J Obstet Gynecol* 1999;181(3):718–724. [PubMed: 10486489]
59. Groenman F, Rutter M, Caniggia I, Tibboel D, Post M. Hypoxia-inducible factors in the first trimester human lung. *J Histochem Cytochem* 2007;55(4):355–363. [PubMed: 17189520]
60. Rodesch F, Simon P, Donner C, Jauniaux E. Oxygen measurements in endometrial and trophoblastic tissues during early pregnancy. *Obstet Gynecol* 1992;80(2):283–285. [PubMed: 1635745]
61. Caniggia I, Mostachfi H, Winter J, et al. Hypoxia-inducible factor-1 mediates the biological effects of oxygen on human trophoblast differentiation through TGFbeta(3). *J Clin Invest* 2000;105(5):577–587. [PubMed: 10712429]
62. De Paepe ME, Mao Q, Powell J, et al. Growth of pulmonary microvasculature in ventilated preterm infants. *Am J Respir Crit Care Med* 2006;173(2):204–211. [PubMed: 16210670]
63. Gappa M, Stocks J, Merkus P. Lung growth and development after preterm birth: further evidence. *Am J Respir Crit Care Med* 2003;168(3):399. author reply 400. [PubMed: 12888614]
64. Greenough A. Late respiratory outcomes after preterm birth. *Early Hum Dev* 2007;83(12):785–788. [PubMed: 17905545]
65. Hjalmarson O, Sandberg K. Abnormal lung function in healthy preterm infants. *Am J Respir Crit Care Med* 2002;165(1):83–87. [PubMed: 11779735]
66. Hoo AF, Dezateux C, Henschen M, Costeloe K, Stocks J. Development of airway function in infancy after preterm delivery. *J Pediatr* 2002;141(5):652–658. [PubMed: 12410193]
67. Jobe AH. An unknown: lung growth and development after very preterm birth. *Am J Respir Crit Care Med* 2002;166(12 pt 1):1529–1530. [PubMed: 12471065]
68. Stocks J, Coates A, Bush A. Lung function in infants and young children with chronic lung disease of infancy: the next steps? *Pediatr Pulmonol* 2007;42(1):3–9. [PubMed: 17123320]
69. Mansell AL, Driscoll JM, James LS. Pulmonary follow-up of moderately low birth weight infants with and without respiratory distress syndrome. *J Pediatr* 1987;110(1):111–115. [PubMed: 3641904]
70. McEvoy C, Bowling S, Williamson K, McGaw P, Durand M. Randomized, double-blinded trial of low-dose dexamethasone: II. Functional residual capacity and pulmonary outcome in very low birth weight infants at risk for bronchopulmonary dysplasia. *Pediatr Pulmonol* 2004;38(1):55–63. [PubMed: 15170874]
71. Morley CJ, Davis PG, Doyle LW, et al. Nasal CPAP or intubation at birth for very preterm infants. *N Engl J Med* 2008;358(7):700–708. [PubMed: 18272893]
72. Friedrich L, Stein RT, Pitrez PM, Corso AL, Jones MH. Reduced lung function in healthy preterm infants in the first months of life. *Am J Respir Crit Care Med* 2006;173(4):442–447. [PubMed: 16322648]
73. Friedrich L, Pitrez PM, Stein RT, Goldani M, Tepper R, Jones MH. Growth rate of lung function in healthy preterm infants. *Am J Respir Crit Care Med* 2007;176(12):1269–1273. [PubMed: 17885265]

74. Stern DA, Morgan WJ, Wright AL, Guerra S, Martinez FD. Poor airway function in early infancy and lung function by age 22 years: a non-selective longitudinal cohort study. *Lancet* 2007;370(9589):758–764. [PubMed: 17765525]
75. Weiss ST, Ware JH. Overview of issues in the longitudinal analysis of respiratory data. *Am J Respir Crit Care Med* 1996;154(6 pt 2):S208–S211. [PubMed: 8970389]
76. Welliver TP, Garofalo RP, Hosakote Y, et al. Severe human lower respiratory tract illness caused by respiratory syncytial virus and influenza virus is characterized by the absence of pulmonary cytotoxic lymphocyte responses. *J Infect Dis* 2007;195(8):1126–1136. [PubMed: 17357048]
77. Raju TN. Late-preterm births: challenges and opportunities. *Pediatrics* 2008;121(2):402–403. [PubMed: 18245431]
78. Raju TN, Higgins RD, Stark AR, Leveno KJ. Optimizing care and outcome for late-preterm (near-term) infants: a summary of the workshop sponsored by the National Institute of Child Health and Human Development. *Pediatrics* 2006;118(3):1207–1214. [PubMed: 16951017]




FIGURE 1.

Proportion of infants with neonatal morbidity as a function of GA. Newborn morbidity was assessed by using a combination of indicators on infants' hospital discharge record and mortality data available from death certificates. Infants born at 34 to 35 weeks' GA were 7 times more likely to have neonatal morbidity than term infants. Infants born at 34 weeks' GA had 20 times the risk of morbidity compared with infants born at 40 weeks' GA. Term (37- to 41-week-GA) infants: $n = 377\ 638$; preterm (34- to 36-week-GA) infants: $n = 26\ 170$. (Adapted with permission from Shapiro-Mendoza CK, Tomashek KM, Kotelchuck M, et al. *Pediatrics*. 2008;121[2]:e227. Available at: www.pediatrics.org/cgi/content/full/121/2/e223.)




FIGURE 2.

A, Correlation between GA and chest-wall compliance. (Reproduced with permission by Blackwell Publishing, Ltd, from Gerhardt T, Bancalari E. *Acta Paediatr Scand.* 1980;69[3]:361.) B, Correlation between age and chest-wall compliance. (Reproduced with permission from Papastamelos C, Panitch HB, England SE, Allen JL. *J Appl Physiol.* 1995;78[1]:182, copyright © 1995 American Physiologic Society.)




FIGURE 3.

Airway-tethering. Tension is transmitted to the airway through the surrounding alveolar septal walls. Airway diameter, and therefore expiratory flows, are increased at higher lung volumes. (Reprinted with permission from Leff A, Schumacker PT. *Respiratory Physiology: Basics and Applications*. Philadelphia, PA: W.B. Saunders Company; 1993:43.)




FIGURE 4.

Growth rate of lung function in healthy preterm infants born at <35 weeks' GA (no significant respiratory disease; patients were excluded if they required any mechanical ventilation, supplemental oxygen for >48 hours, or treatment with surfactant). Shown are flow-volume curves in preterm infants (solid line) versus control term group (dotted line), quantified by the FVC and forced expiratory flow at 75% of FVC (forced expiratory flow [FEF₇₅]) ($P < .05$ between groups). Despite normal lung volume, preterm infants had persistently reduced airflow through the age of 16 months. (Reprinted with permission from Friedrich L, Pitrez PM, Stein RT, Goldani M, Tepper R, Jones MH. *Am J Respir Crit Care Med.* 2007;176[12]:1272, copyright © 2007 American Thoracic Society.)

TABLE 1
 Characteristics of Studies That Evaluated the Epidemiology and Outcomes of Preterm (32–36 Weeks' GA) Birth

Study	Primary Objective	Design and Methods	Comparison Populations	Results	Study Quality or Limitations
General morbidity					
McIntire and Leveno ⁷ (2008)	To analyze neonatal mortality and morbidity rates in 34- to 36-wk-GA and term (39-wk-GA) infants	Retrospective cohort study of all live-born singleton infants of 34–40 wk GA born over a period of 18 y in a center in Dallas, TX	34-wk-GA infants (<i>n</i> = 3498); 35-wk-GA infants (<i>n</i> = 6571); 36-wk-GA infants (<i>n</i> = 11 702); 37-wk-GA infants (<i>n</i> = 26 504); 39-wk-GA infants (<i>n</i> = 84 747; referent)	Neonatal mortality rate per 1000 live births of 1.1, 1.5, and 0.5 at 34, 35, and 36 wk GA, respectively, vs 0.2 at 39 wk GA (<i>P</i> < .001); increased neonatal morbidity rate in 34- to 36-wk-GA infants	Did not examine long-term effects of prematurity; results from this population might not be generalizable to other populations
Melamed et al ²¹ (2009)	To estimate the effect of GA on neonatal morbidity and identify predictors of adverse neonatal outcome	Retrospective study of all spontaneous, low-risk deliveries of 34–36 wk GA from 1997 to 2006 in a center in Tel Aviv, Israel	34- to 36-wk-GA infants (<i>n</i> = 2478); term infants (<i>n</i> = 7434)	Increased risk of neonatal morbidity in 34- to 36-wk-GA infants, including respiratory distress syndrome, sepsis, intraventricular hemorrhage, hypoglycemia, and jaundice requiring phototherapy (all <i>P</i> < .001)	Retrospective design precluded blinding of infant age during analysis; did not examine long-term effects of prematurity
Pulver et al ⁸ (2009)	To compare neonatal and infant mortality rates of late-preterm, early-term, and term newborns; to determine the RR of neonatal and infant death for categories of weight for GA; and to examine causes of neonatal and infant death	Retrospective cohort analysis of birth- and death-certificate data for all infants born in Utah between 1999 and 2005 at ≥34 wk GA	343 322 infants total, categorized as 34–36 wk GA, 37–38 wk GA, 39–42 wk GA, small for GA, appropriate for GA, and large for GA; 9 strata total; appropriate for GA at 39–42 wk GA was the referent	34- to 36-wk-GA small-for-GA infants ~44 times more likely than 39- to 42-wk-GA appropriate-for-GA newborns to die in their first month and ~21 times more likely to die in their first year	Relying on data from birth and death certificates, which could not be corrected if questionable; results from this population might not be generalizable to other populations
Santos et al ⁵ (2008)	To investigate the consequences of term and 34- to 36-wk-GA birth on infant health through 3 mo of age	Population-based birth-cohort study of infants born in Pelotas, Brazil, during 2004	34- to 36-wk-GA infants (<i>n</i> = 447); 37- to 41-wk-GA infants (<i>n</i> = 3262; referent)	The risk of neonatal morbidity was ~3 times higher in 34- to 36-wk-GA infants than 37- to 41-wk-GA infants (25.6% vs 8.1%; <i>P</i> < .001); hospital admission rates in first 3 mo of life were 30% higher in 34- to 36-wk-GA infants; neonatal and infant mortality rates were ~5 times higher in 34- to 36-wk-GA infants	Information on maternal morbidity during pregnancy and on delivery was obtained through maternal recall; the quality of care provided to newborns could not be evaluated
Shapiro-Mendoza et al ²⁶ (2008)	To compare morbidity risk in 34- to 36-wk-GA and 37- to 41-wk-GA infants	Population-based cohort of singleton, 34- to 36-wk-GA and 37- to 41-wk-GA infants who were born in Massachusetts hospitals in Massachusetts	34- to 36-wk-GA infants (<i>n</i> = 26 170); 37- to 41-wk-GA infants (<i>n</i> = 377 638; referent)	34- to 36-wk-GA infants were 7 times more likely to have newborn morbidity than term infants (22% vs 3%); infants born at 34 wk GA had 20 times higher risk of	Other confounding variables may not have been identified because of use of secondary sources; misclassification of medical conditions

Study	Primary Objective	Design and Methods	Comparison Populations	Results	Study Quality or Limitations
Tomashek et al ¹² (2006)	To understand how 34- to 36-wk-GA infants are affected by discharge policies created for 37- to 41-wk-GA infants	Retrospective population-based cohort of singleton, 34- to 36-wk-GA and 37- to 41-wk-GA newborns born to Massachusetts residents in Massachusetts hospitals from January 1, 1998, to November 30, 2002, at a hospital and discharged home early after vaginal delivery	34- to 36-wk-GA infants (<i>n</i> = 1004); 37- to 41-wk-GA infants (<i>n</i> = 24 320; referent)	4.3% 34- to 36-wk-GA vs 2.7% 37- to 41-wk-GA infants were either readmitted or had an observational stay	Hospital readmissions and observational stays could not all be linked to the birth records; unable to account for all neonatal morbidity with data sources used; GA and adequacy of prenatal care were estimated from birth certificates; results from this population might not be generalizable to other populations
Tomashek et al ⁶ (2007)	To assess differences in mortality rate between 34- to 36-wk-GA and 37- to 41-wk-GA infants	Retrospective analysis of data from the US period-linked birth/infant death data for 1995–2002 for singleton 34- to 36-wk-GA and 37- to 41-wk-GA infants born to residents of the 50 states and the District of Columbia	34- to 36-wk-GA infants (<i>n</i> = 2 221 545); 37- to 41-wk-GA infants (<i>n</i> = 24 973 117; referent)	Despite a significant decline in infant mortality rates between 1995 and 2002, infant mortality rates were 3 times higher in 34- to 36-wk-GA infants than 37- to 41-wk-GA infants; early, late, and postneonatal mortality rates were 6, 3, and 2 times higher, respectively, in 34- to 36-wk-GA than in 37- to 41-wk-GA infants	Relied on data from birth and death certificates, which could not be corrected if questionable; cause of death limited to information on death certificate; underlying cause of death may have been misclassified; GA was estimated from birth certificate
Young et al ¹⁰ (2007)	To determine RR for mortality and causes and ages of death in 34- to 42-wk-GA infants	Retrospective cohort analysis of birth- and death-certificate data for all infants born in Utah between 1999 and 2004 at ≥34 to ≤42 wk GA	34-wk-GA infants (<i>n</i> = 3194); 35-wk-GA infants (<i>n</i> = 5722); 36-wk-GA infants (<i>n</i> = 12 190); 39-wk-GA infants (<i>n</i> = 26 175); 40-wk-GA infants (<i>n</i> = 61 764; referent); 41-wk-GA infants (<i>n</i> = 14 361); 42-wk-GA infants (<i>n</i> = 1081)	Most common cause of death in both groups was birth defects; mortality rates were still higher in 34- to 36-wk-GA than in 40-wk-GA infants after excluding those who died of birth defects	GA and causes and ages of death were based on birth- and death-certificate data; results from this population might not be generalizable to other populations; cause for preterm delivery unknown
General respiratory morbidity					
Clark ¹⁷ (2005)	To examine causes of a need for ventilation in near-term infants and to determine rates of mortality, CLD, and neurologic morbidity	Prospective cohort analysis of infants of ≥34 wk GA who were intubated within 72 h and required ventilation for <6 h	Mean GA (SD): 37 wk (2wk) (<i>n</i> = 1011)	51 (5%) patients died; 109 (11%) patients had CLD; 86 (9%) patients had neurologic complications	Small population; patients were selected for the need for a mechanical ventilator

Study	Primary Objective	Design and Methods	Comparison Populations	Results	Study Quality or Limitations
Dani et al ²⁰ (2009)	To compare the rate of clinical complications (hyperbilirubinemia, hypoglycemia, hypothermia, poor feeding, and respiratory distress) in 34- to 36-wk-GA and ≥ 37 -wk-GA infants	Observational population-based study of infants born in 2008 in a center in Florence, Italy, with birth weight of >2000 g and ≥ 34 wk GA	34- to 36-wk-GA infants ($n = 137$); ≥ 37 -wk-GA infants ($n = 2708$; referent)	34- to 36-wk-GA infants had higher rates of clinical complications than ≥ 37 -wk-GA infants	Results from this population might not be generalizable to other populations; only healthy infants were examined
De Luca et al ²⁷ (2009)	To compare mortality rates and incidence of clinically significant morbidities in <37 -wk-GA and ≥ 37 -wk-GA infants according to delivery mode	Prospective analysis of obstetric and neonatal outcomes of all deliveries performed between January 1982 and September 2004 in a center in Geneva, Switzerland	$N = 56\ 549$; neonates were stratified according to GA in strata of 1 wk, from 34–42 completed weeks, and grouped into <37 -wk-GA and ≥ 37 -wk-GA neonates	Depression at birth, admission to special care, and respiratory morbidity showed a strong age-related trend independent of delivery mode; there was an up to >10 -fold decrease in respiratory morbidity from 34 wk GA to term depending on delivery mode	Results from this population might not be generalizable to other populations; restricted indications for elective cesarean delivery may have selected for higher-risk pregnancies
Escobar et al ⁹ (2006)	Quantification of short-term hospital outcomes in 35- to 36-wk-GA infants	Review of published data and reanalysis of databases or retrospective cohort analysis	$n = 47\ 495$; 35- to 36-wk-GA infants; ≥ 37 -wk-GA infants (referent)	Considerable mortality and morbidity in 35- to 36-wk-GA infants; mortality rate = 0.8% in those who survived for 6 h; $\sim 8\%$ required oxygen support for at least 1 h; more likely to be rehospitalized within 14 or 15–182 d	Conclusions were drawn from limited available data
Khashu et al ¹⁹ (2009)	To compare the mortality and morbidity rates in 33- to 36-wk-GA and 37- to 40-wk-GA infants	Population-based study that compared mortality and morbidity data from the British Columbia Perinatal Database Registry of all singleton newborns born at 33–40 wk GA between April 1, 1999, and March 31, 2002	33- to 36-wk-GA infants ($n = 6381$); 37- to 40-wk-GA infants ($n = 88\ 867$; referent)	Stillbirth, perinatal, neonatal, and infant mortality rates were higher in the 33- to 36-wk-GA infants; higher incidence of respiratory morbidity and longer hospital stays in 33- to 36-wk-GA infants	Lack of stratification of certain variables based on underlying diseases; inability to analyze according to week of gestation
Kitsommart et al ²⁸ (2009)	To compare the morbidity and mortality rates of 34- to 36-wk-GA and ≥ 37 -wk-GA infants	Retrospective chart review of 34- to 36-wk-GA infants who were admitted to the NICU, intermediate care nursery (level 2), or postpartum wards at a center in Hamilton, ON, Canada, from April 1, 2004, to March 31, 2008	34- to 36-wk-GA infants ($n = 1481$); ≥ 37 -wk-GA infants ($n = 9332$; referent)	Worse respiratory outcomes in 34- to 36-wk-GA infants than in ≥ 37 -wk-GA infants; respiratory outcomes included rates of positive pressure, ventilation requirement (either continuous positive airway pressure or mechanical ventilation), and prevalence of pneumothorax (all $P < .001$)	Results from this population might not be generalizable to other populations; retrospective design may call into question validity of data
Wang et al ¹³ (2004)	To test whether 35- to 36-wk-GA vs 37- to 40-wk-GA infants have more postnatal	Electronic medical record database sorting and analysis of neonatal records of infants who	35- to 36-wk-GA infants ($n = 90$); 37- to 40-wk-GA infants ($n = 95$; referent)	Similar median length of stay in 35- to 36-wk-GA infants born vaginally or by	Results from this population might not be generalizable to other

Study	Primary Objective	Design and Methods	Comparison Populations	Results	Study Quality or Limitations
Infectious respiratory morbidity					
Boyce et al ²³ (2000)	To determine rates of RSV hospitalization in children with and without specific medical conditions	Retrospective cohort study of all children <3 y old enrolled in the Tennessee Medicaid program from July 1, 1989, to June 30, 1993	Age strata: 0 to <6 mo, 6 to <12 mo, 12 to <24 mo, and 24 to <36 mo; within each stratum, low-risk (referent): BPD, CHD, ≤28 wk GA, 29–33 wk GA, 33–36 wk GA, and other	RSV hospitalization rates were highest in 0-to <6-month-olds and decreased with age; 33- to 36-wk-GA infants had 159.6 hospitalizations per 1000 child-years compared with 163.6 for 29-to <33-wk-GA infants and 88.2 for infants born at term	Database only contains information on children enrolled in Medicaid; hospitalization rates among such children might not be generalizable to the entire population; RSV infection was identified by coding, which may have underestimated RSV impact
Carbonell-Estrany et al ²⁹ (2000)	To collect data on RSV hospitalization rates and to determine risk factors for rehospitalization among premature infants with and without CLD in Spain	Observational, prospective, longitudinal study in 14 Spanish hospitals (annual birth cohort, 57 000 infants) between October 1, 1998, and March 31, 1999	≤28-wk-GA infants (<i>n</i> = 101); 29- to 30-wk-GA infants (<i>n</i> = 179); 31- to 32-wk-GA infants (<i>n</i> = 304); infants with CLD (<i>n</i> = 38)	Lower risk for RSV hospitalization with increasing GA (OR: 0.85 [95% CI: 0.72–0.99]; <i>P</i> < .047); higher risk for hospital admission among infants with CLD (OR: 3.1 [95% CI: 1.22–7.91]; <i>P</i> < .016)	6% of sample lost to follow-up, may have caused underestimation of hospitalization rate; timing of RSV season may have caused underestimation of hospitalization rate; mean sensitivity of RSV test may have underestimated true number of RSV cases
Figueras-Alloy et al ³¹ (2004)	To identify risk factors most likely to lead to RSV-related hospitalization in 33- to 35-wk-GA infants	Hospital-based case-control study conducted in 50 centers in Spain during the RSV season (October 2002–April 2003); controls were selected and included in May–June 2003	33- to 35-wk-GA infants hospitalized for RSV (<i>n</i> = 186); 33- to 35-wk-GA age-matched controls (<i>n</i> = 371; referent)	Risk factors for RSV-related hospitalization were absolute chronological age at start of RSV season; breastfeeding for ≤2 mo; ≥1 school-aged siblings; ≥4 residents in the home; family history of wheezing	Results from this population might not be generalizable to other populations; case-control design may have introduced bias in selection of controls
Holman et al ³⁶ (2003)	To examine the epidemiology of and identify risk factors for bronchiolitis-associated deaths among infants in the United States	Analysis of national multiple cause-of-death and linked birth/infant death data for the years 1995–1998	<i>n</i> = 229 bronchiolitis-associated infant deaths; <32 wk GA; 32–35 wk GA; ≥36 wk GA (referent)	Bronchiolitis-associated mortality rates were 19.7 per 100 000 live births in <32-wk-GA infants, 5.8 in 32- to 35-wk-GA infants, and 1.3 in ≥36-wk-GA infants	Bronchiolitis was identified by coding, which may have underestimated bronchiolitis impact
Horn and Smout ³³ (2003)	To determine if GA is independently associated with hospital-resource use	Analysis of retrospective data for infants hospitalized with RSV bronchiolitis or	≤32-wk-GA infants (<i>n</i> = 28); 33- to 35-wk-GA infants (<i>n</i> = 31); 36-	33- to 35-wk-GA infants had higher resource use than ≤32-wk-GA or ≥37-wk-GA	Data regarding additional risk factors were not collected

Study	Primary Objective	Design and Methods	Comparison Populations	Results	Study Quality or Limitations
Law et al ³⁰ (2004)	and outcomes among infants hospitalized for RSV To identify independent risk factors for hospitalization of infants with RSV infection	pneumonia admitted to 10 children's medical centers from April 1, 1995, to September 30, 1996 Prospective, multicenter, observational cohort study of infants born at 33–35 wk GA between November 1 and ~April 30 of 2000–2001 or 2001–2002 in Canada	wk-GA infants (<i>n</i> = 30); ≥37-wk-GA infants (<i>n</i> = 215) 1832 infants 33–35 wk GA	infants; these differences remained significant in multiple regression analyses 57% (56 of 98) of the infants had an emergency department visit for RSV; 69% (66 of 96) of the infants were hospitalized for RSV	Results from this population might not be generalizable to other populations; relied on culture-confirmed illness; not all patients were tested
Leader and Kohlbase ³² (2003)	To provide current estimates of the incidence, associated risk factors, and costs of severe RSV infections among infants in the United States, defined as emergency department visits, hospitalization, and death	Retrospective analysis of National Hospital Ambulatory Medical Care Survey data (1997–2000), National Hospital Discharge Survey data (1997–2000), and Perinatal Mortality Linked Files (1998–1999)	Children with comorbidities (<i>n</i> = 372); children without comorbidities (<i>n</i> = 288)	Children ≤35 wk GA vs ≥37 wk GA had significantly higher RSV mortality rates (<i>P</i> < .01)	Retrospective design may call into question validity of data
Sampalis ³⁵ (2003)	To evaluate the impact of RSV infections or resource utilization in 32- to 35-wk-GA infants in Canada	Analysis of data from infants born at 32–35 wk GA between 1997 and 2000, hospitalized for proven or probable RSV, and followed to December 31, 2001	32- to 35-wk-GA infants hospitalized for proven or probable RSV (<i>n</i> = 2415); 32- to 35-wk-GA control infants (<i>n</i> = 20 254)	Hospitalization rates, total hospital stay, outpatient visits, and overall deaths highest in 32- to 35-wk-GA infants	Use of database may call into question validity of RSV diagnosis; data regarding other causes that led to use of health care services were not available; direct causality between RSV hospitalization and the antecedent outcomes evaluated was not absolutely certain
Willson et al ³⁴ (2003)	To characterize complications among infants of ≤35 wk GA who were hospitalized for bronchiolitis or RSV pneumonia	Analysis of retrospective data for infants hospitalized with RSV bronchiolitis or pneumonia admitted to 10 children's medical centers from April 1, 1995, to September 30, 1996	≤32-wk-GA infants (<i>n</i> = 44); 33- to 35-wk-GA infants (<i>n</i> = 43); 36-wk-GA infants (<i>n</i> = 46); ≥37-wk-GA infants (<i>n</i> = 338)	Respiratory complications most common (60%) and most serious; highest complication rates, longer stay, and higher costs in 33- to 35-wk-GA infants	Use of database may call into question validity of diagnoses

CLD indicates chronic lung disease; OR, odds ratio; CHD, congenital heart disease.